

The Super-little Higgs

C. Csáki^a, G. Marandella^b, Y. Shirman^c and A. Strumia^d

^a *Institute for High Energy Phenomenology
Newman Laboratory of Elementary Particle Physics
Cornell University, Ithaca, NY, 14853, USA*

^b *Department of Physics, University of California, Davis, CA 95616, USA*

^c *Theoretical Division T-8, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

^d *Dipartimento di Fisica dell'Università di Pisa and INFN, Italia*

csaki@lepp.cornell.edu, maran@physics.ucdavis.edu,
shirman@lanl.gov, Alessandro.Strumia@df.unipi.it

Abstract

Supersymmetry combined with little-Higgs can render the Higgs vev super-little, providing models of electroweak symmetry breaking free from fine-tunings. We discuss the difficulties that arise in implementing this idea and propose one simple successful model. Thanks to appropriately chosen Higgs representations, D -terms give a negligible tree-level mass term to the Goldstone. The fermion representations are anomaly free, generation independent and embeddable into an $SU(6)$ GUT. A simple mechanism provides the large top quark mass.

1 Introduction

Weak-scale supersymmetry (SUSY) is considered the most promising interpretation of the origin of the electroweak symmetry breaking scale. However, the non-discovery of superpartners and the Higgs at LEP and the Tevatron imply that almost all superpartners must be heavier than the W, Z vectors, making typical SUSY models fine-tuned [1]. The essential problem is that the mass parameters in the Higgs potential are determined by the soft SUSY breaking terms. It is then hard to understand how the Higgs VEV and consequently the W, Z masses could naturally be sufficiently smaller than the soft breaking masses themselves. This problem is exacerbated in the Minimal Supersymmetric Standard Model (MSSM) by the fact that one needs significant one loop corrections to the higgs quartic self coupling in order to push the higgs mass above the 115 GeV LEP bound [2]. This can be achieved with heavy and maximally-mixed stops, at the price of more fine-tuning.

Faced with this problem, alternative interpretations have been sought after. One prominent idea [3] is that the Higgs could be a pseudo-Goldstone boson of a global symmetry broken at a scale f . The main problem with this approach is that the Higgs does not in fact behave very much like a Goldstone-boson:

it has a sizable $\mathcal{O}(1)$ quartic self coupling, implying (naively) that any global symmetry should be broken by $\mathcal{O}(1)$ effects to generate such a coupling. This problem has been circumvented in little Higgs models by introducing the idea of collective breaking: the mass of the pseudo-Goldstone can be protected from large corrections (while still allowing for a sizable quartic) if every coupling respects some subset of the global symmetries under which the Goldstone transforms [4, 5]. However, in most little-Higgs models electroweak precision observables are affected at tree level by the new physics introduced to implement this idea, pushing the symmetry breaking scale f in the multi-TeV range [6, 7] such that fine-tuning are needed to render the Higgs vev little enough [8].

One can see that the problems with the two approaches are complementary to each other: supersymmetric models would not give large electroweak precision corrections for soft breaking terms of order a few hundred GeV, however the Higgs potential is fine tuned due to corrections to the soft masses of order $(m_{\text{soft}}/4\pi)^2 \ln \Lambda/m_{\text{soft}}$, where m_{soft} is a typical soft breaking term and Λ is the cutoff scale. With a high cutoff scale $\Lambda \sim 10^{16} - 10^{19}$ GeV the logarithm is basically compensating for the loop factor and so the absence of fine tuning would imply $m_{\text{soft}} \sim M_Z$. In little Higgs models the corrections to the Higgs mass are generically of the form $(f/4\pi)^2 \ln f/v$, where v is the Higgs VEV. This could give a natural Higgs potential for $f \sim \text{few hundred GeV}$, but typically f has to be much larger. If one were able to combine the two approaches, a natural Higgs potential may emerge: with m_{soft} and f are somewhat above current experimental bounds,

$$m_{\text{soft}} \sim \text{few hundred GeV} \quad \text{and} \quad f \sim \text{few TeV},$$

the corrections to the Higgs mass would be given by $(m_{\text{soft}}/4\pi)^2 \log f/m_{\text{soft}}$. See [9, 10, 11] for recent attempts in this direction.

We stress that in this approach the hierarchy problem would be mainly resolved by supersymmetry, and the effect of the pseudo-Goldstone mechanism would be to ensure that the Higgs mass parameter relevant to electroweak symmetry breaking be a loop factor lower than the soft breaking mass parameters (without large logarithmic enhancement) thus naturally making the W, Z bosons lighter than all supersymmetric particles.

When using the lack of positive experimental results as a guideline for exploring new directions one should pay attention in not inventing a medicine that is worse than the illness: a complicated model might look less plausible than the MSSM fine-tuned at a few % level.

In the present case, a very similar mechanism has already been applied to SUSY models when trying to solve the doublet-triplet splitting problem of SUSY GUTs. The perhaps most elegant resolution of this problem is to assume that the Higgs doublets are much lighter than the triplets because they are pseudo-Goldstone bosons. An appealing model of this sort, based on an $SU(6)$ gauge group, has been suggested in [12, 13] improving on [14]. The idea is that the Higgs fields of the model are comprised of two sectors (an $SU(6)$ adjoint Σ and a pair of fundamentals $H, \bar{H}$) which are not directly connected by superpotential terms. Then the Higgs sector has an $SU(6)_{\Sigma} \otimes SU(6)_{H, \bar{H}}$ global symmetry which is broken via the Higgs VEV's around the GUT scale as follows: $SU(6)_{\Sigma} \rightarrow SU(4) \otimes SU(2)_L \otimes U(1)_Y$ and $SU(6)_{H, \bar{H}} \rightarrow SU(5)$. One key feature of the model is that this symmetry-breaking pattern leaves one complex $SU(2)_L$ doublet as Goldstone bosons (and supersymmetry protects a second doublet from a large mass). The other key feature of this model is that while gauging $SU(6)$ explicitly breaks the global symmetry, due to the non-renormalization theorem the Goldstones can not get a supersymmetric mass from the D -terms and pick up a mass only after supersymmetry is softly broken.

The description of the $SU(6)$ GUT model eerily reminds the story of the ‘simplest’ little Higgs model [11], where an $SU(3)_L \otimes U(1)_X$ gauge symmetry is broken by two separate non-communicating sectors leaving two doublets as Goldstones. Since $SU(3)_c \otimes SU(3)_L \otimes U(1)_X \subset SU(6)$ it is most natural to try to apply the particular group theory suggested by the $SU(6)$ model to our attempt at combining supersymmetry with a pseudo-Goldstone mechanism. In fact, it turns out that this approach has some advantages compared to the ‘simplest’ little Higgs:

- Unifying two Higgs doublets in an $SU(3)_L$ adjoint ensures that, in appropriate ranges of parameter space, D -terms do not give dangerous contributions to the Goldstone potential.
- The chiral fermions are automatically anomaly free, generation independent, and embeddable into an $SU(6)$ GUT group.
- A simple mechanism makes the top quark the only SM fermion with a mass comparable to the electroweak scale.
- The gauge couplings may possibly unify. One can find a simple matter content such that the gauge couplings unify at a high scale around M_{Pl} . However, in the full model, the QCD coupling hits a Landau pole around 10^{13} GeV.

This paper is organized as follows. In Section 2 we give a more quantitative description of the fine tuning problems facing supersymmetric and little Higgs models, and a brief summary of our approach to combining supersymmetry with the pseudo-Goldstone mechanism. In Section 3 we present the concrete model we investigate: we describe the gauge, Higgs and matter content, and introduce a top sector, that provides an order one Yukawa coupling only to the top quark. We compute the electroweak precision bounds on the $SU(3)_L$ breaking scales. In Section 4 we calculate the Higgs potential induced by one-loop effects and show how one could generate a tree-level quartic self-interaction term for the Higgs using a version of the sliding singlet mechanism suggested in [11]. In Section 5 we discuss issues related to the phenomenology of the model: we calculate the running of the gauge couplings and comment on possibilities for unification, we discuss the consequences of the presence of extra sterile neutrinos, and comment on possible signatures at the LHC. Finally we conclude in Section 6.

2 Motivation

2.1 The problems of SUSY

Most SUSY models are fine tuned. In order to motivate our extensions of SUSY models it is useful to rephrase this problem in the following way. Let us define f as the RGE energy scale at which EWSB appears: $m_h^2(f) = 0$, where m_h^2 is the running Higgs mass parameter. RGE corrections will drive m_h^2 negative, and in order to have correct EWSB they should be not too large. In models like the constrained minimal supersymmetric standard model (CMSSM), the scale f is dynamically determined by RGE running. Keeping only the dominant top Yukawa contribution to the running of m_h^2 , and assuming large $\tan\beta$ (at tree level this corresponds to a Higgs with mass equal to M_Z) one needs:

$$M_Z^2 = \frac{3y_t^2}{2\pi^2} m_t^2 \ln \frac{f}{m_{\tilde{t}}} \quad (1)$$

where $y_t = m_t/174 \text{ GeV} \approx 1$ and $m_{\tilde{t}}$ is the average stop mass. This contribution to M_Z^2 would generically be much larger than M_Z^2 , and it can be reduced only at the price of fine-tuned cancellations with other contributions. To avoid such fine tunings one needs to impose that this contribution is not unnaturally large. To achieve this one would need

$$m_{\tilde{t}} \lesssim \text{few hundred GeV} \quad \text{and} \quad f \lesssim \text{few TeV}. \quad (2)$$

Models like the CMSSM are problematic because the scale f is dynamically determined by RGE running as $f = M_{\text{GUT}} e^{-F}$ where the function F depends on all dimensionless parameters (like $g_3, \lambda_t, M_{1/2}/m_{\text{soft}}$). Then $f \sim \text{TeV}$ can usually only be obtained thanks to a fine-tuning among these parameters. The main motivation of this paper is to try to find a rationale for this apparent coincidence of f with the electroweak scale.

The simplest possible interpretation is that the Higgs is a pseudo-Goldstone-boson of a global symmetry spontaneously broken at a low energy scale f . Models of this type with $f \sim M_{\text{GUT}}$ have been studied as solutions to the doublet/triplet problem of SUSY GUTs: this would explain why $m_h \ll M_{\text{GUT}}$, but would not make the low-energy Higgs potential more natural. However, recently proposed little-Higgs techniques allow us to lower f down to the TeV scale and thus obtain natural EWSB. This is the direction we will be pursuing in this paper.¹

2.2 The problems of Little Higgs

The Higgs boson does not seem to be the pseudo-Goldstone boson of a global symmetry spontaneously broken at a scale f . Indeed the Goldstone mechanism completely resets the Higgs potential: $V(h) = 0 \cdot |h|^2 + 0 \cdot |h|^4$, while the Higgs must have a substantial quartic coupling. This problem has been recently circumvented by inventing very specific breakings of the global symmetry, that generate a $|h|^4$ term at tree level, while $|h|^2$ only arises at loop level. In models of this kind the generic outcome of pseudo-Goldstone models

$$V(h) \propto \cos^n(|h|/f) \quad \text{with a minimum at } \langle |h| \rangle \sim f \quad (3)$$

gets replaced by $\langle |h| \rangle \sim f y_t / 4\pi$. In the ‘little-Higgs’ limit $\langle |h| \rangle \ll f$ the Goldstone potential can be expanded as

$$V(h) = -m^2 |h|^2 + \lambda |h|^4 + \mathcal{O}\left(\frac{|h|^6}{f^2}\right). \quad (4)$$

The mass term is generated by one-loop corrections as

$$m^2 = 0 \cdot \Lambda^2 + \frac{\mathcal{O}(g^2, \lambda_t^2)}{(4\pi)^2} f^2, \quad (5)$$

where the quadratically divergent piece cancels thanks to the little Higgs mechanism, leaving a logarithmically divergent contribution. Most models assume or need a low cut-off, $\Lambda \lesssim 4\pi f$.

However, the new physics introduced to implement the little-Higgs mechanism, providing the cancellation of the UV-divergent Λ^2 term, typically also gives tree-level corrections to precision data implying $f \gtrsim$ few TeV. As a result a sizable fine-tuning (the typical one corresponding to the ‘little hierarchy problem’ [17]) is needed to render m^2 small enough. This situation is so bad because the actual gain achieved by the little-Higgs mechanism is more modest than what is suggested by the naïve estimate $\langle |h| \rangle / f \sim y_t / 4\pi$: the top loop happens to be multiplied by big $\mathcal{O}(1)$ coefficients. Indeed without a tree level quartic one would get $\langle |h| \rangle / f \sim 1$, and the top loop correction to the quartic is almost enough to get the desired higgs mass, $m_h \gtrsim 115 \text{ GeV}$ [2]: adding a tree level quartic does not make the higgs as little as desired, $\langle |h| \rangle \ll f$. One needs to make m^2 smaller than its natural value given by Eq. (5), by invoking fine-tuned cancellation between its contributions.

If we were to combine supersymmetry with the little Higgs mechanism then SUSY will naturally lower m^2 , giving

$$m^2 \sim \frac{\mathcal{O}(g^2, \lambda_t^2)}{(4\pi)^2} m_{\text{soft}}^2 \ln \frac{f}{m_{\text{soft}}}. \quad (6)$$

A super-little (slittle) higgs $\langle |h| \rangle \ll f$ is now obtained for $m_{\text{soft}} \ll f$.

2.3 SUSY and little-Higgs: a difficult marriage

We discussed why supersymmetric little-Higgs models are an interesting possibility. We now discuss the obstacles that must be overcome to actually implement this idea. The main incompatibility is that supersymmetry suppresses λ more than m^2 , providing a natural mechanism to do exactly the opposite of what the little-Higgs mechanism is supposed to do. More concretely:

¹See [15, 16] for alternative tentative interpretations of the smallness of f .

- Supersymmetric D -terms break the global symmetry already at tree level, making the soft pseudo-Goldstone mass term not smaller than soft terms of other sparticles.
- Supersymmetry forbids superpotential contributions to the quartic Higgs coupling, if the low energy theory has the MSSM field content.

The second problem can be overcome by building little-Higgs models that provide at tree level a NMSSM-like $SH\bar{H}$ coupling, where S is a singlet. The most pressing problem is the first one. One can hope that a simple solution exists, because D -terms are not a generic breaking of the global symmetry, but they have a very specific structure. Indeed the reason why D -terms are present in supersymmetric theories is the following: when a gauge symmetry is broken, D -terms give mass to the scalar superpartner of the Goldstone equal to the vector boson mass, thereby completing a SUSY massive vector multiplet. Therefore it is worth investigating if the specific structure of the D -terms can be used to bypass the first problem.

Let us start looking at the ‘simplest’ model [10, 11]. The basic idea consists in promoting $SU(2)_L \otimes U(1)_Y$ to $SU(3)_L \otimes U(1)_X$, and breaking $SU(3)_L$ by two pairs of Higgs superfield triplets, $(H_1, \bar{H}_1)$ and $(H_2, \bar{H}_2)$. In the absence of couplings between the two pairs, the potential has an $SU(3)_{H_1\bar{H}_1} \otimes SU(3)_{H_2\bar{H}_2}$ symmetry. Since it is partly gauged, one of the two Goldstone bosons is eaten by the $SU(3)_L/SU(2)_L$ vector bosons, while the orthogonal combination remains a pseudo-Goldstone boson. Such pseudo-Goldstone field G is contained in the triplets as

$$H_1 = e^{i\Pi F_2/F_1 F}(0, 0, f_1/\sqrt{2}), \quad \bar{H}_1 = (0, 0, \bar{f}_1/\sqrt{2})e^{-i\Pi F_2/F_1 F} \quad (7)$$

$$H_2 = e^{i\Pi F_1/F_2 F}(0, 0, f_2/\sqrt{2}), \quad \bar{H}_2 = (0, 0, \bar{f}_2/\sqrt{2})e^{-i\Pi F_1/F_2 F} \quad (8)$$

where $F_i^2 = (f_i^2 + \bar{f}_i^2)/2$, $F^2 = F_1^2 + F_2^2$ and Π is the Goldstone 3×3 matrix

$$\Pi = \begin{pmatrix} 0_2 & G \\ G^\dagger & 0 \end{pmatrix}.$$

The part of the D -term potential that breaks the $SU(3)_{H_1\bar{H}_1} \otimes SU(3)_{H_2\bar{H}_2}$ global symmetry is:

$$V_D = \frac{g^2}{8} \left(|H_1^\dagger \cdot H_2|^2 - |\bar{H}_1 \cdot H_2|^2 - |\bar{H}_2 \cdot H_1|^2 + |\bar{H}_2^\dagger \cdot \bar{H}_1|^2 \right) + \dots \quad (9)$$

and its contribution to the Goldstone potential is

$$V_D = \frac{g^2}{8} (f_1^2 - \bar{f}_1^2)(f_2^2 - \bar{f}_2^2) \cos^2 \left[\frac{\sqrt{G^\dagger G}}{F} \left(\frac{F_1}{F_2} - \frac{F_2}{F_1} \right) \right] + \dots \quad (10)$$

The vevs $f_i, \bar{f}_i$ have to be computed by minimizing the full potential, such that in general they depend on G .² We see that Goldstones typically get a mass comparable to other sparticles, because H_i and $\bar{H}_i$ typically have different soft masses (the difference is induced e.g. by top Yukawa renormalization effects) leading to VEVs that are non-supersymmetric: $f_i^2 - \bar{f}_i^2 \sim m_i^2 - \bar{m}_i^2$. A generic supersymmetric little-Higgs model does not provide the desired super-little Higgs vev.

On the other hand, if the VEVs were supersymmetric, $\langle H_1 \rangle = \langle \bar{H}_1 \rangle$ and $\langle H_2 \rangle = \langle \bar{H}_2 \rangle$ then no potential is generated along the Goldstone direction. One way of naturally forcing the VEVs to be supersymmetric is by implementing a symmetry between H_i and $\bar{H}_i$ that keeps the soft terms equal: $m_{H_i}^2 = m_{\bar{H}_i}^2$ [11], or by

²When one of the two sectors is heavy and supersymmetric, this reproduces at low energy the usual MSSM quartic Higgs couplings, such that a Goldstone quartic potential arises if $\tan\beta \neq 1$. In this model $\tan\beta$ is determined by the vev of an extra light field, present because the global symmetry has an extra $U(1)$ factor, absent in the model that will be considered in this paper.

	$SU(3)_c$	$SU(3)_L$	$U(1)_X$
H	1	3	+1/3
$\bar{H}$	1	$\bar{3}$	-1/3
Σ	1	8	0
$2 \times D_{1,2,3}$	$\bar{3}$	1	+1/3
$2 \times L_{1,2,3}$	1	$\bar{3}$	-1/3
$U_{1,2,3}$	$\bar{3}$	1	-2/3
$E_{1,2,3}$	1	$\bar{3}$	+2/3
$Q_{1,2,3}$	3	3	0
Q'	3	3	-1/3
$\bar{Q}'$	3	$\bar{3}$	+1/3

Table 1: *Charge assignments of the chiral superfields. The upper group is the Higgs sector, the middle group is the chiral matter, the lower group is the vector-like top.*

building an appropriate top sector where only one Higgs couple is involved in large Yukawa couplings [10]. Indeed eq. (10) shows that to preserve the Goldstone flat direction it is enough to have supersymmetric VEVs only in one couple: either $f_1 = \bar{f}_1$ or $f_2 = \bar{f}_2$. This consideration suggests that it might be sufficient to simply replace one of the two doublet/anti-doublet pairs with an $SU(3)_L$ adjoint Σ , that contains a doublet/anti-doublet Higgs pair: their soft masses are automatically forced to be equal by $SU(3)_L$ gauge-invariance, and one expects that this is enough for avoiding the unwanted D -term potential for the Goldstones. (Replacing both doublets with adjoints would not allow to get the top mass). Models of this type with $f \sim M_{\text{GUT}}$ have already been explored as an elegant solution to the doublet/triplet splitting problem [12], and as a source for the top mass [13]. By ‘deconstructing’ $SU(6)$ into $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ fragments gives a model analogous to the ‘simplest’ supersymmetric little-Higgs [10, 11], and with universal family charges, no gauge anomalies, and a source for the top mass.

In conclusion, we are lead to study the model described in detail in the next section.

3 The model

3.1 The gauge sector

We consider an $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ supersymmetric gauge theory. The Higgs sector, as discussed in the previous section, consists of one $SU(3)_L$ triplet and anti-triplet pair, H and $\bar{H}$, and an adjoint Σ .

Unlike in previous attempts [10, 11, 18, 19, 20], our assignment of fermion gauge quantum numbers, listed in Table 1, is anomaly free, generation universal and reproduces the SM chiral content. To see easily that these goals are indeed achieved we observe that our gauge group can be embedded into $SU(6)$, which contains one generation of SM fermions in a $15 \oplus \bar{6} \oplus \bar{6}'$ representation. Indeed, one can immediately check that it decomposes under the usual $SU(5)$ gauge group as $10_{15} \oplus \bar{5}_{15} \oplus \bar{5}_{\bar{6}} \oplus 1_{\bar{6}} \oplus \bar{5}'_{\bar{6}} \oplus 1'_{\bar{6}'}$, so that the chiral part is the usual $10 + \bar{5}$. Decomposing it into representations of our $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ group gives the charge assignment shown in the first two parts of Table 1.

The superpotential of the Higgs sector of our model is³

$$W_{\text{Higgs}} = \frac{M}{2} \text{Tr} \Sigma^2 + \frac{\lambda}{3} \text{Tr} \Sigma^3 + S(\lambda'' H \bar{H} - M'^2) + \lambda' \bar{H} \Sigma H \quad (11)$$

³We normalize the super-fields such that their kinetic terms are $\int d^4\theta [2\text{Tr} \Sigma^\dagger \Sigma + H^\dagger H + \bar{H}^\dagger \bar{H}]$.

The dimensionful M, M' terms are analogous to the MSSM μ term. The $SU(3)_L$ -breaking Higgs VEVs are

$$\langle \Sigma \rangle = \text{diag}(w/2, w/2, -w), \quad \langle H \rangle = (0, 0, f/\sqrt{2}), \quad \langle \bar{H} \rangle = (0, 0, \bar{f}/\sqrt{2}) \quad (12)$$

and break the gauge group to the SM gauge group, giving the following masses to the extra gauge bosons, a complex weak doublet W' and a weak singlet Z' :

$$M_{W'}^2 = \frac{g^2}{2}(F^2 + 9w^2), \quad M_{Z'}^2 = \frac{2F^2 g^2}{3 - t^2} \quad (13)$$

where $F^2 = (f^2 + \bar{f}^2)/2$ and $t = \tan \theta_W$. The unbroken $U(1)_Y$ arises as $Y = X + T_8/\sqrt{3}$ (on triplets $T_8/\sqrt{3} = \text{diag}(1/6, 1/6, -1/3)$). The hypercharge gauge coupling is $1/g'^2 = 1/g_X^2 + 1/3g^2$, and g is the usual $SU(2)_L$ coupling.

We will need VEVs an order of magnitude larger than the SUSY-breaking scale. They can arise because M, M' are larger than the SUSY-breaking scale, or because the λ, λ'' couplings are small. The model can work even with a vanishing M .

For $\lambda' = 0$ the superpotential decomposes as $W(H, \bar{H}) + W(\Sigma)$, acquiring a $SU(3)_{H, \bar{H}} \otimes SU(3)_\Sigma$ global symmetry. The Higgs sector consists of 4 complex doublets, out of which one is eaten during the breaking of $SU(3)_L$ and one acquires a heavy mass from the D -terms, forming a massive supersymmetric vector multiplet with the corresponding gauge bosons. We are then left with two complex doublets, H_u, H_d which are contained in the $SU(3)_L$ multiplets as follows

$$H = \exp\left(i\Pi \frac{3w}{FV}\right) \langle H \rangle, \quad (14a)$$

$$\bar{H} = \langle \bar{H} \rangle \exp\left(-i\Pi \frac{3w}{FV}\right) \quad (14b)$$

$$\Sigma = \exp\left(-i\Pi \frac{F}{3wV}\right) \langle \Sigma \rangle \exp\left(i\Pi \frac{F}{3wV}\right) \quad (14c)$$

where $F^2 \equiv (f^2 + \bar{f}^2)/2$, $V^2 \equiv F^2 + 9w^2$, the VEVs are defined in Eq. (12) and Π is a 3×3 matrix defined as

$$\Pi = \frac{1}{\sqrt{2}} \begin{pmatrix} 0_2 & H_u \\ H_d^t & 0 \end{pmatrix}. \quad (15)$$

Only the combination $G \equiv (H_u + H_d^\dagger)/\sqrt{2}$ is a Goldstone boson, and this is the field we want to identify the Higgs with. The orthogonal sGoldstone $\tilde{G} \equiv (H_u - H_d^\dagger)/\sqrt{2}$ is massless as long as supersymmetry is unbroken and gets a mass of the order of the soft susy breaking mass for the Higgs.

The λ' term explicitly breaks the global symmetry giving a tree level mass to the Goldstone. Since a successful model does not require λ' to be smaller than a ‘typical’ SM Yukawa coupling — $\lambda' \lesssim 10^{-2}$ is small enough — it seems unnecessary to invent a mechanism that suppresses λ' . From this point of view, the situation is simpler than in the GUT-scale version of the model.

Gauge and top Yukawa interactions break the $SU(3)_{H, \bar{H}} \otimes SU(3)_\Sigma$ global symmetry, generating a mass and a quartic coupling for the pseudo-Goldstone G .

3.2 The top sector

As usual in supersymmetric theories we will assume the presence of matter parity: a Z_2 symmetry under which ‘matter’ chiral superfields flip sign, while the fields in the Higgs sector are even. Or we could impose R -parity, under which also the superspace coordinate θ_α flips sign. These will forbid B, L violating operators.

In little Higgs models one can try to impose a different Z_2 symmetry, usually referred to as T -parity [21], in order to forbid tree-level corrections to electroweak precision observables. In our case this would correspond to imposing every new particle beyond the SM to be odd: this is neither necessary nor possible. It is not possible because this symmetry is not compatible with the $SU(3)_L \otimes U(1)_X$ gauge group. It is not necessary because we do not need to lower the little-Higgs scale F, w below a few TeV.

After imposing this symmetry, the most general renormalizable superpotential term involving the matter fields is

$$W_{\text{matter}} = \alpha_{ij} Q_i \bar{H} D_j + \beta_{ij} E_i \bar{H} L_j. \quad (16)$$

The VEV of $\bar{H}$ gives a mass term that mixes the extra set of D, L fields with $SU(3)_L$ partners of SM fields. Thus below the little-Higgs scale the matter content of the theory is exactly that of the MSSM, plus possibly two sterile neutrinos per generation from the third components of the L fields. We will comment on these sterile neutrinos later on. One can also see that at the renormalizable level (16) does not lead to any mass terms for the MSSM fields even after $SU(2)_L$ breaking. Perhaps the simplest way to see this is by performing an $SU(3)_L$ rotation on the $\bar{H}$ field such that its VEV (after $SU(2)_L$ breaking) is still contained in the third component. Then all of the $SU(2)_L$ breaking would be rotated into the Σ field, which however does not participate in (16). Thus at this level all the MSSM fermions remain massless.

Therefore in order to implement the little-Higgs mechanism we must find a way of obtaining a top Yukawa coupling compatible with the gauge symmetries of our model. This can be elegantly achieved following the mechanism [13] employed in the $SU(6)$ model: by adding the appropriate vector-like representation listed in the lower rows of Table 1, additional Yukawa couplings are allowed, such that one up-type quark gets a mass $m_t \sim v$. These extra fields form a heavy vector-like top analogous to the one employed by little-Higgs models, but different from the one employed in the closely related ‘simplest’ little Higgs.

Let us first review the details of this mechanism in the $SU(6)$ theory. We add the representation 20 (three index antisymmetric tensor) to the other matter fields $3 \times (15 + \bar{6} + \bar{6}')$. Then the most general renormalizable superpotential is

$$W = 20 \Sigma 20 + 20 H 15 + 15 \bar{H} \bar{6} \quad (17)$$

where here $H, \bar{H}, \Sigma$ denote the obvious $SU(6)$ -extensions of our $H, \bar{H}, \Sigma$ fields. The 20 representation is pseudo-real: despite being non-chiral, $SU(6)$ invariance keeps it massless. The way the large Yukawa coupling is generated in the $SU(6)$ model is by assuming that the VEV of $H, \bar{H}$ break the symmetry to $SU(5)$. Under this subgroup

$$20 = 10_{20} + \bar{10}_{20}, \quad 15 = 10_{15} + 5_{15}, \quad \bar{6} = \bar{5}_{\bar{6}} + 1_{\bar{6}}, \quad \bar{6}' = \bar{5}'_{\bar{6}'} + 1'_{\bar{6}'}.$$

When $H, \bar{H}$ get a VEV, the Yukawa coupling $20 H 15$ gives a $\bar{10}_{20} 10_{20}$ mass term, and $15 \bar{H} \bar{6}$ gives a $5_{15} \bar{5}_{\bar{6}}$ mass term. The light chiral fermions arise as $10_{20} \oplus \bar{5}'_{\bar{6}'}$. The first term of eq. (17) provides the top Yukawa coupling to the Higgs Goldstone in Σ . This mechanism has the collective symmetry breaking already built in.

Implementing this mechanism in the $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ model corresponds to adding the representations $Q' = (\bar{3}, 3)_{-1/3}$ and $\bar{Q}' = (3, \bar{3})_{1/3}$ (which make up the 20 of $SU(6)$ up to two singlets). Therefore, in terms of the $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ fragments the top sector is given by

$$W_{\text{top}} = \lambda_1 \bar{Q}' \Sigma Q' + \lambda_2 \bar{Q}' H U + \lambda_3 Q H Q' \quad (18)$$

The top sector contains the SM top quark and 3 additional heavy vector-like particles. It has the collective breaking property that any individual coupling respects the global $SU(3)_\Sigma \otimes SU(3)_H$ symmetry, broken only by the simultaneous presence of both λ_1 and λ_2 (or λ_1 and λ_3), thus eliminating the quadratically divergent loop effect from the top sector.

A $\bar{Q}' Q'$ mass term is also allowed, and should be not much bigger than $\lambda_1 \langle \Sigma \rangle$. Its presence would modify the detailed expressions for the masses and for the loop induced potential calculated below, however

qualitatively does not change the story: the symmetry breaking is still collective. In the $SU(6)$ unified theory this mass term is not allowed, and it remains vanishing if the 20 does not couple to the $SU(6)$ -breaking Higgses.

The top mass matrix is given by

$$W_{\text{top}} = (\bar{Q}', Q) M_{\text{top}} (Q', U)^t \quad M_{\text{top}} = \begin{pmatrix} \lambda_1 \Sigma & \lambda_2 H \\ \lambda_3 \Omega & 0 \end{pmatrix}$$

where $\Omega_{ij} = \epsilon_{ijk} H_k$ is a 3×3 matrix. To find the spectrum we diagonalize the matrix $M_{\text{top}}^\dagger M_{\text{top}}$. Neglecting $SU(2)_L$ breaking, one gets a massless mode (to be identified with the SM top) and three heavy modes ('heavy tops') with masses

$$M_{T_1}^2 = \lambda_1^2 w^2 + \frac{1}{2} \lambda_2^2 f^2, \quad M_{T_{2,3}}^2 = \frac{1}{4} (\lambda_1^2 w^2 + 2 \lambda_3^2 f^2). \quad (19)$$

The SM top Yukawa coupling y_t arises in terms of $\lambda_{1,2,3}$ as

$$y_t = \frac{f^2 \sqrt{F^2 + 9w^2} \lambda_1 \lambda_2 \lambda_3}{F \sqrt{2(2w^2 \lambda_1^2 + f^2 \lambda_2^2)(w^2 \lambda_1^2 + 2f^2 \lambda_3^2)}} \simeq \begin{cases} \lambda_1/2 & \text{if } w \rightarrow 0 \\ 3f^2 \lambda_2 \lambda_3 / (2Fw \lambda_1) & \text{if } F \rightarrow 0 \end{cases}. \quad (20)$$

At this stage only the top is massive, while all other Standard Model fermions remain massless. Their masses can arise from non-renormalizable superpotential operators in a way similar to the construction of [13], where fermion masses are suppressed by powers of $M_{\text{GUT}}/M_{\text{Pl}}$. In our model the natural suppression is given by $\epsilon = f/\Lambda'$, where Λ' is some scale associated with the explicit breaking of the global $SU(3)_{H,\bar{H}} \otimes SU(3)_\Sigma$ symmetry. We will see in Section 4.2 that such a scale is necessarily present in order to generate a tree-level quartic self coupling for the Higgs, so it is natural to assume that the same scale will appear in these non-renormalizable operators (most of which will break $SU(3)_{H,\bar{H}} \otimes SU(3)_\Sigma$).

3.3 Electroweak precision constraints

Precision data are affected at tree level in two different ways: by the Z' , and by the presence of a Higgs triplet coupled to the Higgs doublet. We can neglect the extra corrections induced at loop level by supersymmetric particles and by the heavy vector-like tops.

The effects of the Z' have already been studied in [7], and data imply the bound $F > 3 \text{ TeV}$ at 99% C.L. The Higgs triplet effects can be computed from eq. (14c): expanding it in the limit $H_u, H_d \ll w \ll F$ one finds that the $SU(2)_L$ adjoint contained in Σ gets a VEV $\langle \Sigma \rangle^{ij} = H_d^i H_u^j / 12w$. Assuming that $SU(2)_L$ is dominantly broken by the Goldstone VEV, such that $\langle H_u \rangle = \langle H_d \rangle = (0, v/\sqrt{2})$, one gets a correction $\hat{T} = v^2/144w^2$ to the $\hat{T}$ parameter (normalized as in [30]). Data roughly demand $\hat{T} \lesssim 10^{-3}$ [30] implying $w \gtrsim 0.5 \text{ TeV}$.

4 The Higgs potential and electroweak symmetry breaking

We now study if the motivations discussed above can be realized in the present model, making sparticles naturally heavier than present experimental bounds.

To start, one can verify that the D -terms potential

$$V_D \propto V_D(H, \bar{H}) + V_D(\Sigma) - \bar{H}[\Sigma, \Sigma^\dagger] \bar{H}^\dagger + H^\dagger[\Sigma, \Sigma^\dagger] H \quad (21)$$

despite breaking at tree level the $SU(3)_{H,\bar{H}} \otimes SU(3)_\Sigma$ global symmetry, does not give any contribution to the Goldstone potential: $|G|^2$ does not appear in $[\Sigma, \Sigma^\dagger]$. However, the full potential have to be minimized with respect to heavy fields: in the supersymmetric limit this eliminates the D -terms corresponding to the

broken generators. For example, in the limit $f \gg w, m_H, m_{\bar{H}}$, the low-energy theory below the scales $f, \bar{f}$ only contains the $SU(2) \otimes U(1)$ D -terms, plus soft terms of order $(m_H^2 - m_{\bar{H}}^2)D_8$ [23, 10], where $m_H, m_{\bar{H}}$ are the soft breaking terms for $H, \bar{H}$, and D_8 denotes the D -term corresponding to the T_8 generator of $SU(3)$ with only the light fields included. This produces a negligibly small Goldstone mass, of order

$$\frac{(m_H^2 - m_{\bar{H}}^2)(f^2 - \bar{f}^2)w^2}{(f^2 + \bar{f}^2)^2}|G|^2. \quad (22)$$

The fact that no large mass for the Goldstone is generated by the D -terms is a necessary but not a sufficient condition, because it does not guarantee that the Goldstone Higgs doublet plays the main rôle in electroweak symmetry breaking. The main rôle would be played by non-Goldstone Higgs fields, if their squared mass parameters are negative (or more precisely if the full squared Higgs mass matrix has a negative eigenvalue).

Let us therefore study the full Higgs potential. At energies below F, w the little-Higgs sector decouples leaving the MSSM with specific values of its Higgs parameters. At tree level and for $\lambda' = 0$ there is a flat Goldstone direction: $H_u = H_d$ i.e. $\tan \beta = 1$. Therefore the Higgs mass matrix, in the basis $(H_u, H_d^\dagger)$ has the form [22]

$$\begin{pmatrix} m_{h_u}^2 + \mu^2 & -B\mu \\ -B\mu & m_{h_d}^2 + \mu^2 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} m_G^2 + m_{G\tilde{G}}^2 & -m_{\tilde{G}}^2 \\ -m_{\tilde{G}}^2 & m_{\tilde{G}}^2 - m_{G\tilde{G}}^2 \end{pmatrix} \quad (23)$$

where m_G^2 and $m_{G\tilde{G}}^2$ are given in terms of high-energy parameters as:

$$m_G^2 = \frac{F^2}{V^2}m_\Sigma^2 + \frac{9w^2}{F^2V^2}(m_H^2f^2 + m_{\bar{H}}^2\bar{f}^2), \quad m_{G\tilde{G}}^2 \sim m_H^2 - m_{\bar{H}}^2. \quad (24)$$

where $m_\Sigma, m_H, m_{\bar{H}}$ denote soft mass terms. We only showed the most important feature of the full expression of $m_{G\tilde{G}}^2$. The Goldstone/sGoldstone mixing $m_{G\tilde{G}}^2$ tends to induce a VEV along the $\tan \beta \neq 1$ direction. In this way a large enough Higgs mass can be achieved, but the motivation for the model is lost: the Higgs is no longer a pseudo-Goldstone boson and the low energy theory is neither better nor worse than the generic MSSM.

This problem does not arise in a well defined region of the parameter space. Choosing $m_{\tilde{G}} \sim m_\Sigma \sim \text{TeV}$ a few times larger than $m_H, m_{\bar{H}}$ suppresses the unwanted negative eigenvalue $-m_{G\tilde{G}}^4/4m_{\tilde{G}}^2$ of eq. (5). Furthermore, $m_{G\tilde{G}}^2$ can be naturally small: with w a few times smaller than f , the top Yukawa coupling is dominantly generated by λ_1 : λ_2, λ_3 can be small enough such that RGE running does not break an initial equality $m_H^2 = m_{\bar{H}}^2$. The initial equality naturally arises in models of soft terms (like gauge mediation and possibly gravity mediation) where the source of supersymmetry breaking couples equally to H and $\bar{H}$. Furthermore, we assume $m_G^2 > 0$. This is a plausible assumption, since our model does not need many Yukawa couplings as large as the top Yukawa, that renormalize soft squared masses making them negative.

4.1 The top loop effect

At this point we can turn on the loop effects that break the global symmetry, and study if Goldstones acquire a small $SU(2)_L$ -breaking VEV. One loop corrections renormalize the parameters in eq. (23) and more importantly generate a potential for the Goldstone mode. Expanding the potential along the Goldstone direction as $V(G) = -m^2|G|^2 + \lambda|G|^4$, it receives at one loop order two main contributions: from the sector that provides the top Yukawa, and from gauge interactions. The first contribution is

$$\lambda \simeq \frac{3y_t^4}{8\pi^2} \left(\ln \frac{m_{\tilde{t}}}{m_t} + \frac{A_t^2}{2m_{\tilde{t}}^2} \left(1 - \frac{A_t^2}{12m_{\tilde{t}}^2} \right) \right) \quad (25a)$$

$$m^2 \simeq \frac{3}{8\pi^2} y_t^2 (m_{\tilde{t}}^2 - m_t^2) \left(2 \ln \frac{M_{T_2}}{m_{\tilde{t}}} + 1 + 2c \ln \frac{M_{T_1}}{M_{T_2}} \right) \quad (25b)$$

where we assumed a common stop mass $m_{\tilde{t}}$, A_t is the top A -term, and

$$c = \frac{f^2 (F^2 + 9w^2) \lambda_1^2 (2w^2 (\lambda_2^2 - \lambda_3^2) \lambda_1^2 + f^2 \lambda_2^2 (\lambda_2^2 - 4\lambda_3^2))}{6F^2 (2w^2 \lambda_1^2 + f^2 \lambda_2^2) (2 (\lambda_2^2 - \lambda_3^2) f^2 + 3w^2 \lambda_1^2) y_t^2} \quad (26)$$

As expected λ depends only on MSSM physics and takes its standard value, and m^2 depends only on physics up to $\mathcal{O}(M_{T_{1,2}})$ energies: it is generated by one-loop threshold corrections at the scale $M_{T_{1,2}}$, and by short RGE running down to m_t , in agreement with the estimate in eq. (6).

Adding also the gauge contribution and making m^2 a few times smaller by appropriately chosen gaugino masses, the Goldstone potential can develop a minimum. However, even in the most favorable situation of maximal stop mixing $A_t = \sqrt{6}m_{\tilde{t}}$, to get $m_h \gtrsim 115$ GeV one needs $\tan \beta \gtrsim 3$ i.e. the model is allowed only when its motivation is lost.

4.2 A tree level quartic

To reach the goal outlined in the introduction, we must introduce a little-Higgs like mechanism that provides an extra tree-level contribution to the quartic potential. This can be done analogously to [11].

We assume that the low energy theory contain extra singlet(s) S , such that a NMSSM-like SG^2 superpotential term is allowed. However, it is not easy to obtain only this term from the original theory. For example breaking the $SU(3)_{H,\bar{H}} \otimes SU(3)_\Sigma$ global symmetry by adding terms like $S\bar{H}\Sigma H$ would also generate a Goldstone mass term of order F, w . Indeed, assuming for simplicity $\bar{f} = f$,

$$\bar{H}\Sigma H = -\frac{f^2 w}{2} + \frac{V^2}{12w} |G|^2 \quad (27)$$

and so $|\bar{H}\Sigma H|^2$ contains an unwanted mass term of order f^2, w^2 for the Goldstone.

One needs operators that contain only the Goldstone fields, without the constant VEVs f, w . One can eliminate these VEVs by adding extra singlets, that “slide” absorbing the $\Sigma, H\bar{H}$ VEVs, analogously to the sliding singlet mechanism [24] for the doublet-triplet splitting problem of SUSY GUTs. Naively, this could be achieved by an operator of the form $S(S' + \bar{H}\Sigma H)$. The S equation of motion fixes $\langle S' \rangle = -\langle \bar{H}\Sigma H \rangle$, eliminating the $\Sigma, H, \bar{H}$ VEVs, and thus leaving the Goldstone massless. However S' eliminates also the full dependence on the Goldstone, and thus no potential is generated for G . To be successful, we need a “collective version” of this sliding singlet mechanism, whereby only the VEVs get absorbed, while Goldstone interactions survive. One successful operator is

$$\frac{1}{\Lambda} (SH + S' \frac{\Sigma H}{\Lambda'}) (\bar{S}\bar{H} + \Sigma \bar{H}). \quad (28)$$

Indeed the S and S' equations of motion are respectively

$$\begin{aligned} \bar{S}\bar{H}H + \bar{H}\Sigma H &= 0, \\ \bar{S}\bar{H}\Sigma H + \bar{H}\Sigma^2 H &= 0. \end{aligned} \quad (29)$$

Expanding these equations at leading order in the Goldstone field gives

$$\begin{aligned} \bar{S}f^2 + (-wf^2 + V^2|G|^2/6w) &= 0, \\ \bar{S}(-wf^2 + V^2|G|^2/6w) + (f^2w^2 - V^2|G|^2/6) &= 0. \end{aligned} \quad (30)$$

The Goldstone appears differently in the two equations, while the VEVs appear in the same way: $\bar{S}$ ‘slides’ to $\langle \bar{S} \rangle = w$ eliminating only the VEVs, such that the resulting potential does not contain the unwanted $|G|^2$ mass term, but contains the desired $|G|^4$ coupling with $\mathcal{O}(1)$ coefficient. The peculiar operator in eq. (28) could be naturally generated by tree-level exchange of an extra heavy pair of $SU(3)_L$ triplets.

Another simple mechanism for obtaining a tree-level quartic has been suggested in [10], that exploited the fact that, if one of the Higgs sectors have large VEVs and universal soft breaking terms, the effective theory below the Higgs VEV only contains the D -terms of the unbroken gauge generators, but with $SU(3)$ -symmetric soft-terms. Since the MSSM D -terms do not give a mass for the Goldstone, this offers the possibility of using the usual MSSM quartic for $\tan\beta > 1$. In our model this would correspond to considering the limit $w \gg f$, such that the effective theory below w contains the $SU(2)_L \otimes U(1)_X \otimes U(1)_8$ D -terms (the broken D -terms are eliminated in the process of integrating out the heavy fields). However, the $SU(2)_L \otimes U(1)_X \otimes U(1)_8$ D -terms give the potential

$$-\frac{g^2(f^2 - \bar{f}^2)^2}{4(f^2 + \bar{f}^2)}|G|^2 + \frac{2g^2(f^2 - \bar{f}^2)^2}{3(f^2 + \bar{f}^2)^2}|G|^4 \quad (31)$$

for the Goldstone G . Thus the quartic is sizeable only if f and $\bar{f}$ deviate $\mathcal{O}(1)$ from each other, but in this case a large Goldstone mass is also introduced. To implement this mechanism in our model, one would have to introduce another pair of triplets $\tilde{H}, \bar{\tilde{H}}$ with large VEV's and $\mathcal{O}(1)$ couplings to Σ (in order to avoid the appearance of another global $SU(3)$ symmetry). With these additional fields the low-energy effective theory only contains the $SU(2) \otimes U(1)$ D -terms, that provide only the quartic (but no mass term) for the Goldstone

$$\frac{g^2(f^2 - \bar{f}^2)^2}{8(f^2 + \bar{f}^2)^2}|G|^4 \quad (32)$$

which is the usual MSSM quartic at $\tan\beta > 1$. The value of $\tan\beta$ is given by the ratio $f/\bar{f}$, which is mainly determined by the soft breaking terms for $H, \bar{H}$, and in this limit can be made different from one without introducing large corrections to the Goldstone mass. In the limit $w \gg f, \bar{f}$, by choosing an appropriately small λ_1 and appropriately large λ_2, λ_3 our top sector provides a large top Yukawa coupling $y_t \leq 3f\lambda_2\lambda_3/\sqrt{2}F(\lambda_2 + 2\lambda_3)$.

5 Phenomenology

5.1 Gauge coupling unification?

We now turn to the question of the unification of the gauge couplings. We recall that in order to achieve an anomaly free matter content we have chosen to embed the gauge group of our model $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ into a GUT $SU(6)$ gauge group with the Standard Model fermions transforming in the $15 \oplus \bar{6} \oplus \bar{6}'$ representation. Furthermore, we have added an extra chiral fields in the representation 20 of $SU(6)$ in order to obtain the large top quark mass compatibly with the little Higgs mechanism. It is therefore natural to ask whether unification of couplings could be achieved with an appropriate embedding of the Higgs sector into $SU(6)$. In order to find unification, we can complete the H and $\bar{H}$ multiplet into 6 and $\bar{6}$ representations of $SU(6)$ and assume that the color triplets contained in these fields obtain their mass at a low scale. Finally, Σ can be embedded into the adjoint of the $SU(6)$ gauge group: we will assume that all additional components of Σ have large masses — this constitutes an analog of the doublet-triplet splitting problem.

With the matter content described above we can now calculate RG evolution of the gauge couplings. The RGE β -function coefficient above the SUSY-breaking and the little-Higgs scale are (1, 4, 10) for the $SU(3)_c$, $SU(3)_L$ and $U(1)_X$ groups respectively. We here assumed the $SU(6)$ -unified normalization of the $U(1)_X$ charges, in which the $H, \bar{H}$ fields have charges $\pm 1/\sqrt{6}$. We have not included the fields $Q, \bar{Q}'$ introduced for generating the top mass. One can see in fig. 1a that the gauge couplings can unify at a scale somewhat higher than the usual SUSY $SU(5)$ scale. Taking into account the full 20 of $SU(6)$ does not affect unification, since adding a complete $SU(6)$ multiplet does not change the differential running nor the unification scale. (To generate the top Yukawa we only needed the 18 $Q', \bar{Q}'$ fields; the 20 contains two extra singlets with X -charges ± 1). Fig. 1b shows the running in the minimal model, with a light 20 and without light Higgs




Figure 1: *One-loop running of the $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ gauge couplings in a model with 3 light generations (a) without the matter fields added to generate the top mass and with light colored Higgs triplets: SU(6) unification can happen at a scale somewhat above the usual SU(5) GUT scale; (b) with the matter fields added to generate the top mass and Higgs and without light colored Higgs triplets: a dedicated non-perturbative analysis would be needed to test if unification happens.*

colored triplets: the QCD coupling hits a Landau pole at energies around 10^{13} GeV.⁴ However, this does not automatically imply that there is something wrong. Techniques developed in the past years allow us to analyze strongly interacting supersymmetric gauge theories. The appearance of the QCD Landau pole implies that the description valid at low energies breaks down at these intermediate energies. At higher energies the theory can be described by a dual weakly-coupled gauge group [26]. From a high-energy point of view, the QCD gluons are composite particles, which appear only at low energies as the effective description of some more fundamental gauge theory. The possibility that the SM gauge bosons are composites has been discussed in [26]. In our case, the gauge group above the QCD Landau pole would be $SU(10) \otimes SU(3)_L \otimes U(1)_X$. As it usually happens with product gauge groups, now the $SU(10)$ is asymptotically free. At slightly higher energies the $SU(3)_L$ gauge coupling hits its own Landau pole. Thus we obtain a sort of a duality cascade, that is reminiscent of the Klebanov-Strassler model [27]. There it has been conjectured that the duality cascades at high energies would correspond to a string model on a warped throat, with the supersymmetric SM appearing only at low energies. In the MSSM all gauge couplings remain perturbative up to the Planck scale, so this conjectured cascading behavior seems not well motivated. On the contrary, it can arise in the model at hand. It would be interesting to find out if the apparent unification of the couplings at non-perturbative values will leave any consequences behind for the cascading theory.

5.2 The extra fermions

In our model fermions have a generation-universal and anomaly-free assignment of gauge charges. The basic difference with respect to [10, 11] is that singlet leptons are extended to $SU(3)_L$ triplets rather than to $SU(3)_L$ singlets.

Let us discuss the phenomenology of the extra fermions. Each generation contains an extra vector-like

⁴The minimal model gives successful perturbative SU(6) unification if SUSY is split, that is if only the Higgses have light superpartners. We ignore this possibility because it does not explain the hierarchy problem, unless one resorts to anthropic arguments [25]. Alternatively, perturbative $SU(6)^2$ unification can be achieved if duplicated copies of the $SU(3)_c, SU(3)_L, U(1)_X$ gauge groups survive unbroken down to low enough energies. Here we prefer to insist on minimality.

pair of right-handed down-type quarks with mass $\sim f$ (contained in D, Q), an extra vector-like pair of left-handed leptons with mass $\sim f$ (contained in E, L), and two light $SU(2)_L$ singlets (contained in the two L), that we will name ‘sterile neutrinos’.

The heavy extra fermions decay into ordinary fermions via $SU(3)_L$ gauge interactions. The sterile neutrinos have a more interesting phenomenology. They are light because renormalizable couplings do not allow any $\mathcal{O}(f)$ mass term. A similar phenomenon was observed in [28]. Unlike ordinary sterile neutrinos, the singlets can be produced at accelerators thanks to their $SU(3)_L$ interactions.

In cosmology, these $g = 12$ degrees of freedom of sterile neutrinos remain in thermal equilibrium down to $T_{\text{dec}} \approx \text{GeV} \cdot (f/20 \text{ TeV})^{4/3}$. At this temperature the number of SM degrees of freedom is uncertain, and varies by about a factor 2 depending on whether T_{dec} is above or below QCD phase transition occurring at imprecisely known temperature $T \sim m_\pi$. We assume $g_{\text{SM}}(T_{\text{dec}}) \sim 50$, such that sterile states have temperature $T_s = (10.75/g_{\text{SM}}(T_{\text{dec}}))^{1/3} T_\nu$. Until relativistic, sterile states provide an extra energy density that is conventionally expressed in terms of ‘extra number of neutrinos’ as $\Delta N_\nu \sim 1.5$. This excess is still considered acceptable [29, 31], and can be tested by CMB and BBN experiments.

One can modify the model such that sterile states get $\mathcal{O}(f)$ masses (thereby avoiding phenomenological problems or signatures), by adding more singlets that can get Dirac masses with the steriles. Such extra singlets are automatically present in the full $SU(6)$ model. However, analogously to the usual see-saw, it is considered more plausible that extra singlets also have Majorana masses much larger than f, w . Integrating them out results into $SU(3)_L$ -invariant Majorana neutrino mass operators, that now involve both active and sterile neutrinos: $(LH)^2 \sim (\nu_L v + \nu_s f)^2$. The presence of two L fields per generation prevents a univocal prediction for the sterile neutrino masses in terms of active neutrino masses. Nevertheless, a typical pattern emerges: sterile neutrinos with masses $m_{\nu_s} \sim (f/v)^2 m_\nu \sim 10 \text{ eV}$ and active/sterile mixings $v/f \sim 0.03$. These masses and mixings are dangerously close to the border of the presently excluded region [29], and possibly lie in the range suggested by the ‘sterile’ interpretation of the LSND anomaly [32].

5.3 Consequences for the LHC

The model presented here predicts various sets of new particles, that could be observable at the LHC.

First of all, there are the ‘usual’ supersymmetric particles. The stops are expected to be around a few hundred GeV. Depending on the SUSY-breaking model, this typically implies that most sparticles lie in this same energy range. Usual R -parity conservation is expected in this model, so the traditional search methods for supersymmetry should be applicable.

Next, at the little-Higgs scale $f \sim \text{few TeV}$ there should be a well-defined set of extra gauge bosons, Higgses, fermions, and in particular the vector-like tops. All these new particles are accompanied by their own supersymmetric partners.

Concerning gauge bosons, we expect a heavy $SU(2)_L$ doublet W' and a neutral Z' with masses in the few TeV range. The lower bounds on these masses from electroweak precision data are $M_{W'} \gtrsim 1.5 \text{ TeV}$ and $M_{Z'} \gtrsim 1.7 \text{ TeV}$. Searching for W', Z' in collisions of SM fermions is analogous to searching for W, Z in $e, \bar{e}$ collisions. The Z' couples to pairs of SM fermions, giving a resonant contribution to four-fermion processes: thanks to the clean signature and to the relatively large cross section, the LHC sensitivity is in the multi-TeV range. This issue has been studied for typical Z' bosons [33, 34]. The model univocally predicts all Z' couplings.

Discovering the W' is more difficult, because it couples one SM fermion to one exotic fermion (a heavy quark and lepton, or a light sterile) such that W' ’s need to be pair-produced, giving no resonance and rates more limited by the center-of-mass energy available at the LHC. Single production of W' is also possible, but only thanks to couplings between the W' and two SM fermions suppressed by v^2/f^2 [33].

The vector-like top quarks could have masses in the $2 \div 3 \text{ TeV}$ range. The LHC reach for this kind of particles is expected to be around $2 \div 2.5 \text{ TeV}$ [33, 34]. Not being odd under R -parity, heavy tops can be singly produced.

The $SU(3)_L$ partners of the light fermions are similarly expected to have multi-TeV masses. The LHC reach for these extra fermions is expected to be around a few TeV.

6 Conclusions

Supersymmetry and little-Higgs are respectively characterized by the scales m_{soft} and f . Present experimental bounds

$$m_{\text{soft}} \gtrsim \text{few hundred GeV} \quad \text{and} \quad f \gtrsim \text{few TeV} \quad (33)$$

imply that typical supersymmetric models and typical little-Higgs models are fine-tuned, but that supersymmetric little-Higgs models can still naturally explain electroweak-symmetry breaking, provided that the Higgs mass parameter arises as

$$m_h^2 \sim \frac{\mathcal{O}(g^2, \lambda_t^2)}{(4\pi)^2} m_{\text{soft}}^2 \ln \frac{f}{m_{\text{soft}}} \quad (34)$$

such that the Higgs gets a super-little VEV v . This is however not the generic outcome of supersymmetric little-Higgs models. Rather, D -terms typically break at tree level the global symmetry such that the Goldstone Higgs mass is not suppressed by a loop factor, and furthermore non-Goldstone Higgses can dominantly contribute to $SU(2)_L$ breaking.

Therefore the goal consists in finding a non-generic and possibly simple supersymmetric little-Higgs model that achieves Eq. (34). Doublet/triplet splitting models based on the $SU(6)$ unification group already contain almost all necessary ingredients, and can operate at low energies $f \sim \text{few TeV}$ provided that $SU(6)$ breaks as

$$SU(6) \xrightarrow{M_{\text{GUT}}} SU(3)_c \otimes SU(3)_L \otimes U(1)_X \xrightarrow{f} SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \xrightarrow{v} SU(3)_c \otimes U(1)_{\text{em}}$$

rather than following the conventional $SU(5)$ route. $SU(3)_L$ is broken independently to the SM gauge group by a pair of Higgs triplets $H, \bar{H}$ and by an adjoint Σ , that contains another pair of Higgs doublets. The resulting pseudo-Goldstone is light enough provided that the superpotential coupling $\bar{H}\Sigma H$ has a coefficient smaller than $\sim 10^{-2}$. The little-Higgs structure is analogous to the ‘simplest’ little-Higgs, with various advantages. The model has an anomaly-free, generation-independent and unified fermion content and a ‘collective’ source for the top Yukawa coupling. Furthermore, thanks to the adjoint, in a well defined range of parameter space the D -term problem simply does not arise.

Electroweak symmetry breaking is then triggered by the negative one-loop contribution of the top sector to the mass squared of the Goldstone. Little-Higgs model building is needed to satisfy the Higgs mass bound, $m_h > 115 \text{ GeV}$, adding an order one quartic self interaction among the Goldstones. This was achieved using a collective version of the sliding singlet mechanism.

Gauge couplings can unify in the presence of appropriate simple $SU(6)$ -embeddable light chiral superfields. However, in the full model, gauge couplings hit a Landau pole below the unification scale, leading to a cascading gauge theory that could directly unify into string theory on a warped throat. The fermion sector contains light sterile neutrinos with superweak $SU(3)_L$ interactions and masses f/v times higher than neutrino masses. The LHC should be able to detect (beyond the ordinary MSSM superpartners) at least the Z' and hopefully some of the color triplet top partners as well.

Acknowledgments We thank Z. Berezhiani, M. Cirelli, A. Falkowski, T. Gregoire, M. Perelstein, S. Pokorski, R. Rattazzi and J. Terning for useful discussions. C.C., G.M., and Y.S. would also like to thank the Aspen Center for Physics, where part of this work was done, for its hospitality. The research of C.C. is supported in part by the DOE OJI grant DE-FG02-01ER41206 and in part by the NSF grants PHY-0139738 and PHY-0098631. The work of G.M. is supported in part by Department of Energy grant DE-FG02-91ER40674. The research of Y.S. is supported by the U.S. Department of Energy under contract

References

- [1] Fine-tuning was defined in R. Barbieri and G. F. Giudice, Nucl. Phys. B 306, 63 (1988). The impact of LEP2 negative searches was discussed in A. Strumia, hep-ph/9904247.
- [2] R. Barate *et al.* [ALEPH Collaboration], Phys. Lett. B 565, 61 (2003) [hep-ex/0306033].
- [3] D. B. Kaplan and H. Georgi, Phys. Lett. B 136, 183 (1984).
- [4] N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Lett. B 513, 232 (2001) [hep-ph/0105239].
- [5] N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, JHEP 0207, 034 (2002) [hep-ph/0206021]; N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, T. Gregoire and J. G. Wacker, JHEP 0208, 021 (2002) [hep-ph/0206020]. For a review and complete list of references see M. Schmaltz and D. Tucker-Smith, hep-ph/0502182.
- [6] C. Csáki, J. Hubisz, G. D. Kribs, P. Meade and J. Terning, Phys. Rev. D 67, 115002 (2003) [hep-ph/0211124]; Phys. Rev. D 68, 035009 (2003) [hep-ph/0303236]; J. L. Hewett, F. J. Petriello and T. G. Rizzo, JHEP 0310, 062 (2003) [hep-ph/0211218]; Z. Han and W. Skiba, Phys. Rev. D 72, 035005 (2005) [hep-ph/0506206].
- [7] G. Marandella, C. Schappacher and A. Strumia, Phys. Rev. D 72, 035014 (2005) [hep-ph/0502096].
- [8] For a recent complete analysis of fine-tuning issues see J. A. Casas, J. R. Espinosa and I. Hidalgo, JHEP 0503 (2005) 038 [hep-ph/0502066]; F. Bazzocchi, M. Fabbrichesi and M. Piai, arXiv:hep-ph/0506175.
- [9] P. H. Chankowski, A. Falkowski, S. Pokorski and J. Wagner, Phys. Lett. B 598, 252 (2004) [hep-ph/0407242]; A. Birkedal, Z. Chacko and M. K. Gaillard, JHEP 0410, 036 (2004) [hep-ph/0404197].
- [10] Z. Berezhiani, P. H. Chankowski, A. Falkowski and S. Pokorski, hep-ph/0509311.
- [11] T. Roy and M. Schmaltz, hep-ph/0509357.
- [12] Z. G. Berezhiani and G. R. Dvali, Bull. Lebedev Phys. Inst. 5 (1989) 55 [Kratk. Soobshch. Fiz. 5 (1989) 42]. R. Barbieri, G. R. Dvali and M. Moretti, Phys. Lett. B 312, 137 (1993). Z. Berezhiani, Phys. Lett. B 355 (1995) 481 [hep-ph/9503366]. Z. Berezhiani, C. Csáki and L. Randall, Nucl. Phys. B 444 (1995) 61 [hep-ph/9501336]. C. Csáki and L. Randall, Nucl. Phys. B 466 (1996) 41 [hep-ph/9512278].
- [13] R. Barbieri *et al.*, Nucl. Phys. B 432, 49 (1994) [hep-ph/9405428].
- [14] K. Inoue, A. Kakuto and H. Takano, Prog. Theor. Phys. 75, 664 (1986); A. A. Anselm and A. A. Johansen, Phys. Lett. B 200, 331 (1988); R. Barbieri, G. R. Dvali and A. Strumia, Nucl. Phys. B 391, 487 (1993);
- [15] The mechanism suggested in R. Barbieri and A. Strumia, Phys. Lett. B 490, 247 (2000) [hep-ph/0005203], assumes that the solution to the cosmological constant problems behaves in an appropriate way. This possibility could be implemented in the context of string anthropic interpretations of the cosmological constant problem as discussed by R. Rattazzi at the HEP2005 conference (www.lip.pt/events/2005/hep2005).
- [16] See e.g. K. Choi, K. S. Jeong, T. Kobayashi and K. i. Okumura, hep-ph/0508029; R. Kitano and Y. Nomura, hep-ph/0509039.
- [17] R. Barbieri and A. Strumia, hep-ph/0007265.
- [18] F. Pisano, V. Pleitez Phys. Rev. D 46 (1992) 410; P. H. Frampton, Phys. Rev. Lett. 69, 2889 (1992).
- [19] M. Schmaltz, JHEP 0408, 056 (2004).
- [20] O. C. W. Kong, Phys. Rev. D 70 (2004) 075021.
- [21] H. C. Cheng and I. Low, JHEP 0408, 061 (2004).
- [22] R. Rattazzi, Phys. Lett. B 375 (1996) 181.
- [23] Y. Kawamura, H. Murayama and M. Yamaguchi, Phys. Rev. D 51, 1337 (1995) [hep-ph/9406245].
- [24] E. Witten, Phys. Lett. B 105, 267 (1981); D. V. Nanopoulos and K. Tamvakis, Phys. Lett. B 113, 151 (1982).
- [25] V. Agrawal, S. M. Barr, J. F. Donoghue and D. Seckel, Phys. Rev. D 57 (1998) 5480 [hep-ph/9707380].
- [26] N. Seiberg, Nucl. Phys. B 435, 129 (1995).
- [27] I. R. Klebanov and M. J. Strassler, JHEP 0008, 052 (2000) [hep-th/0007191].
- [28] A. G. Dias, C. A. de S. Pires and P. S. Rodrigues da Silva, Phys. Lett. B 628, 85 (2005) [hep-ph/0508186].
- [29] For a recent complete analysis see M. Cirelli, G. Marandella, A. Strumia and F. Vissani, Nucl. Phys. B 708 (2005) 215 [hep-ph/0403158].
- [30] R. Barbieri, A. Pomarol, R. Rattazzi and A. Strumia, Nucl. Phys. B 703 (2004) 127 [hep-ph/0405040].
- [31] R. H. Cyburt, B. D. Fields, K. A. Olive and E. Skillman, Astropart. Phys. 23 (2005) 313 [astro-ph/0408033].
- [32] LSND collaboration, Phys. Rev. Lett. 81 (1998) 1774 [nucl-ex/9709006]. LSND collaboration, Phys. Rev. D 64 (2001) 112007. [hep-ex/0104049].
- [33] T. Han, H. E. Logan, L. T. Wang, hep-ph/0506313.
- [34] G. Azuelos *et al.*, Eur. Phys. J. C 39S2, 13 (2005).